



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 295/12, 333/60, 333/54 A61K 31/16, 31/33, 31/34 A61K 31/38	A1	(11) International Publication Number: WO 93/19056 (43) International Publication Date: 30 September 1993 (30.09.93)
(21) International Application Number: PCT/CA93/00124 (22) International Filing Date: 26 March 1993 (26.03.93) (30) Priority data: 858,060 26 March 1992 (26.03.92) US (71) Applicant: THE UNIVERSITY OF BRITISH COLUMBIA [CA/CA]; 6328 Memorial Road, Vancouver, British Columbia V6R 1Z3 (CA). (72) Inventors: MacLEOD, Bernard, A. ; 4036 West 13th Avenue, Vancouver, British Columbia V6R 2T4 (CA). WALKER, Michael, J., A. ; 5176 Connaught Drive, Vancouver, British Columbia V6M 3G3 (CA). WALL, Richard, A. ; 3181 West 24th Avenue, Vancouver, British Columbia V6L 1R7 (CA).		(74) Agents: NASSIF, Omar, A. et al.; McCarthy Tétrault, Suite 4700, Toronto Dominion Bank Tower, Toronto-Dominion Centre, Toronto, Ontario M5K 1E6 (CA). (81) Designated States: AU, BR, CA, CZ, FI, HU, JP, KR, NO, NZ, PL, RU, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: AMINOCYCLOHEXYLAMIDES FOR ANTIARRHYTHMIC AND ANAESTHETIC USES (57) Abstract This invention provides aminocyclohexylamides for use within methods of blocking sodium channels in cardiac or neuronal tissue and for use in the manufacture of medicaments to block sodium channels in cardiac or neuronal tissue. This invention also provides commercial kits comprising a pharmaceutical composition containing aminocyclohexylamides and, instructions for the use of the pharmaceutical composition for arrhythmia or for the inducement of local anaesthesia.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

DescriptionAMINOCYCLOHEXYLAMIDES FOR ANTIARRHYTHMIC
AND ANAESTHETIC USES

5

Technical Field

This invention is generally directed toward aminocyclohexylamide compounds that block cardiac and neuronal sodium channels for use in cardiac arrhythmias and the inducement of local anaesthesia, respectively, and for use in the manufacture of medicaments to block sodium channels in cardiac or neuronal tissue.

15 Background of the Invention

Compounds that block sodium channels in the membranes of cardiac or neuronal tissues are known. For example, Class I antiarrhythmic compounds, such as lidocaine, will block membrane channels for sodium ions in cardiac tissue. A subset of such antiarrhythmic compounds (known as Class Ia antiarrhythmics) includes quinidine and procainamide which are capable of blocking potassium channels in cardiac tissue in addition to blocking sodium channels. Such antiarrhythmic compounds will also block sodium channels in neuronal tissue which is a property of a local anaesthetic compound.

Class I antiarrhythmic compounds may be used to treat supraventricular arrhythmias and ventricular arrhythmias. Treatment of ventricular arrhythmia is very important since such an arrhythmia, especially ventricular fibrillation, can be fatal. Serious ventricular arrhythmias (ventricular tachycardia and ventricular fibrillation) occur most often in the presence of myocardial ischemia and/or infarction. Ventricular fibrillation often occurs in the setting of acute myocardial ischemia, before infarction fully develops. At present, lidocaine is the current drug of choice for

prevention of ventricular fibrillation. However, many Class I antiarrhythmic compounds may actually increase mortality in patients who have had a myocardial infarction. Therefore, there is a need in the art to
5 identify new antiarrhythmic treatments, particularly treatments for ventricular arrhythmias. The present invention fills the need, and further provides other related advantages.

It has been suggested that opioid antagonists,
10 such as naloxone or opioid agonists, may interact with sodium channels and have antiarrhythmic activity. However, the opioid activity is probably separate from any such antiarrhythmic activity, as the former activity appears to be stereospecific but the latter activity is
15 not (Sarne, Y., et al. (1991) *Brit. J. Pharmacol.*).

International patent application WO 86/07257 published December 18, 1986, suggested an antiarrhythmic utility for certain aminocycloalkylamide compounds previously known to be analgesics. One such compound is
20 known as U-50,488H and has been shown to be an opioid agonist particularly active at the kappa receptor. (Von Voightlander, P.F., et al. (1983) *J. Pharmacol. Exp. Ther.*, 244:7-12). It has also been suggested that U-50,488H may have a local anaesthetic activity by
25 reducing sodium conductance (Alzheimer, C. and Ten Bruggencate, G. (1990) *J. Pharmacol. Exp. Ther.*, 255:900-905). The analgesic and local anaesthetic properties of U-50,488H are antagonized by the kappa opioid antagonist, naloxone. However, other researchers
30 have been unable to demonstrate any significant antiarrhythmic activity of U-50,488H (Sitsapesan, R., and Parratt, J.R. (1989) *Br. J. Pharmacol.*, 97:795-800) and, it has been predicted that U-50,488H is actually arrhythmogenic (Wong, T.M., et al. (1990) *J. Mol. Cell*
35 *Cardiol.*, 22:1167-1175).

A different group of analgesic aminocyclohexylamide compounds are described in Horwell's

United States Patent Nos. 4,579,863; 4,598,087; 4,656,182; 4,663,343; 4,737,493; and 4,855,316. The opioid activity of these compounds is antagonized by naloxone and is stereospecific with respect to the enantiomers at the amine and amide substituents of the cyclohexyl ring (see: Meecham, K.G., et al. (1989) *Eur. J. Pharmacol.*, 173:151-157; Singh, L. et al. (1990) *Eur. J. Pharmacol.*, 191:477-480; and Hunter, J.C., et al. (1990) *Br. J. Pharmacol.*, 101:183-189).


10

Summary of the Invention

Briefly stated, the present invention provides an enantiomer or geometric isomer of a compound of formula I, or a pharmaceutically acceptable salt thereof, for use within a method of blocking sodium channels in cardiac or neuronal tissue.

15

In a compound of formula I:




(I)

20

n is either 0 or 1; R₁, R₂, R₃, R₄ are hydrogen, hydroxy, alkoxy or from one to four carbon atoms, or points of attachment of a spiro- or fused five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; R₅ and R₆ are either alkyl of from one to five carbon atoms or, when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl, piperidinyl, or hexahydroazepinyl ring; and Q is selected from the group of substituents comprising: 3,4,5-trimethylphenoxy;

25


4



(II)


where R_7 is hydrogen, fluorine, chlorine, alkyl of from one to six carbon atoms, or aryl; Z is $-CH_2-$, $-O-$, $-S-$, or

5 $N-R_8$ where R_8 is hydrogen, alkanoyl of from one to six carbon atoms, or alkyl of from one to six carbon atoms;



(III)


10 where R_9 and R_{10} are independently hydrogen, fluorine, bromine, alkyl of from one to six carbon atoms, or alkoxy of from one to four carbon atoms;



(IV)

15

where R_9 and R_{10} are defined as above; and



(V)

20 where R_{11} and R_{12} are independently hydrogen, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl.

The present invention is also directed toward use of an enantiomer or geometric isomer of a compound of formula I as described above, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use in blocking sodium channels in cardiac or neuronal tissue.

Within a related aspect, the present invention also provides a commercial kit comprising a pharmaceutical composition containing at least one compound of formula I as described above, or a pharmaceutically acceptable salt thereof; and, instructions for the use of the pharmaceutical composition for arrhythmia or to induce local anaesthesia.

Detailed Description of the Invention

The disclosure of the present invention shows that a group of aminocyclohexylamide compounds, including those described in certain United States patents to Horwell, will block sodium channels in cardiac and neuronal tissue. This activity appears to be independent of the opioid agonist activity of the compounds, as the sodium blocking activity is not antagonized by naloxone. This activity is very potent in respect of cardiac tissue and, thus, the compounds are useful as antiarrhythmic agents. As compared to known antiarrhythmic agents, these compounds are particularly potent, effective and fast acting. At higher concentrations, the compounds also block sodium channels locally in neuronal tissue and have a quick onset of action which results in the compounds being useful as local anaesthetic agents. In addition, since the ability of the compounds to block sodium channels is not stereospecific, enantiomers of the compounds may be selected that have the desired sodium blocking activity, but have significantly reduced kappa agonist activity.

Compounds of formula I constitute a class of substituted aminocyclohexamide compounds. One nitrogen atom is an amine nitrogen substituted with R₅ and R₆ as defined above. Preferably, R₅ is methyl and R₆ is a lower alkyl, most preferably methyl, or, when taken together with the nitrogen atom to which they are attached, R₅ and R₆ preferably form a pyrrolidinyl ring or a hexahydroazepinyl ring. The other nitrogen atom is an N-methylamide substituted as described above, wherein n is preferably 1.

Preferably R₁ and R₄ are hydrogen or, R₃ and R₄ are hydrogen and R₁ and R₂ are an oxaspiran ring.

As used herein, the term "aryl" means phenyl; phenyl substituted with alkyl of from one to four carbon atoms, alkoxy of from one to four carbon atoms, nitro, or trifluoromethyl; 2- or 3-thienyl; and, 2- or 3-thienyl substituted with alkyl of from one to four carbon atoms or alkoxy of from one to four carbon atoms.

In formula II depicted above, the bond that links the substituent with the remainder of the compound of formula I is shown as intersecting both rings of the fused ring structure of the substituent II. This indicates that the bond may be at any one of the carbon atoms in the fused ring structure except at the position of R₇.

While a compound of the present invention is depicted above in formula I by a structural formula having a particular isomeric form, such structural formula contains one or more asymmetric carbon atoms and therefore exists in various stereoisomeric forms. In addition, the compound is capable of existing in different geometric isomeric forms. For example, the substituent R₁ of the cyclohexane ring may be positioned on the same side of the average plane of the ring as the amide nitrogen, or on the side opposite. The present invention contemplates the use of all geometric and stereoisomeric forms of the compounds of formula I.

Compounds of formula I may be used in the present invention as individual enantiomers, as racemic mixtures or, as combinations of different substituted compounds of formula I. Examples of individual enantiomers include compounds 2 and 7 below. Examples of racemic mixtures include compounds 1 and 6 below.

The following compounds of formula I are preferred:

1. (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
2. (1R,2R)-(+) -N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
3. [(±)-(1α,2β,4β,5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
4. [(±)-(1α,2β,4β,5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
5. (±)-trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
6. (±)-trans-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;
7. [5R(5α,7α,8β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;
8. (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
9. (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-3-acetamide;
10. [5S(5α,7α,8β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;
11. (1S,2S)-2-(benzo[b]thiophen-4-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
12. (1R,2R)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
13. (1S,2S)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
14. (1R,2R)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
15. (1S,2S)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

16. (1R,2R)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
17. (1S,2S)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
- 5 18. [1S(1 α ,2 β ,4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide;
19. [1R(1 α ,2 β ,4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide;
20. (1R,2R)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide; and
- 10 21. (1S,2S)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide.

The compounds of formula I may be prepared by
15 known methods, including those described in the
aforementioned United States patents to Horwell (which are
incorporated herein in their entirety by reference).
Suitable methods for the synthesis of diaminocyclohexane
intermediates useful for preparation of a variety of
20 compounds identified above are described in Szmuszkovicz,
J., and Von Voightlander, P.F. (1982) *J. Med. Chem.*,
25:1125-1126. The oxaspiro and methoxy-cyclohexanediamine
intermediates useful for syntheses of compounds 7, 10, 18,
and 19 are described in Halfpenny, P.R., et al. (1990) *J.*
25 *Med. Chem.*, 33:286-291. Preparation or sources of the
carboxylic acids used in the final stage of the syntheses
of the compounds listed above are also to be found in the
above references as well as in Clark, C.R., et al. (1988)
J. Med. Chem., 31:831-836. The above latter three
30 references contain information on all the steps of the
syntheses of the compounds listed above, and provide
sufficient guidance to a person skilled in the art to
repeat the synthesis, isolation, and purification of these
and many other analogous compounds. The individual
35 enantiomers are obtained, if desired, from mixtures of the
different forms by known methods of resolution, such as
the formation of diastereomers, followed by
recrystallisation.

The following compounds are particularly preferred as antiarrhythmics:

- (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
- 5 (1R,2R)-(+) -N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
- [(±)-(1 α ,2 β ,4 β ,5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
- 10 [(±)-(1 α ,2 β ,4 β ,5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide; and
- (±)-trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide.
- 15

It is most preferable that the compounds of formula I selected for use in the present invention be those enantiomers that lack kappa opioid activity or wherein such activity is significantly reduced. For this

20 purpose, the stereoisomeric form illustrated in formula I above is preferred. The following enantiomers having reduced kappa opioid activity are preferred:

- [5S(5 α ,7 α ,8 β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-benzo[b]furan-4-acetamide;
- 25 (1R,2R)-2-(benzo[b]thiophen-4-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
- (1R,2R)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
- 30 (1R,2R)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
- (1R,2R)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
- [1R(5 α ,2 β ,4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide; and
- 35 (1R,2R)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide.

The compounds of formula I may be in the form of a pharmaceutically acceptable acid addition salt. Such

salts include the hydrochloride, sulfate, phosphate, citrate, and other salts known in the art. Pharmaceutical compositions of compound I or salts of compound I may contain pharmaceutically acceptable carriers, which are
5 well known in the art.

In order to assess whether a compound has the required pharmacological activity within the present invention, it is subjected to a series of tests. In the first of such tests, a compound is given as increasing
10 (doubling with each dose) intravenous boluses every 8 minutes to a pentobarbital anesthetized rat. The effects of the compound on blood pressure, heart rate and the ECG are measured 30 seconds and 8 minutes after each dose. Increasing doses are given until the animal dies. The
15 cause of death is identified as being of respiratory or cardiac origin. This test gives an indication as to whether the compound is blocking sodium channels and/or potassium channels, and in addition gives information about acute toxicity. The indices of sodium channel
20 blockade are increasing P-R interval and QRS widening of the ECG. Potassium channel blockade results in Q-T interval prolongation of the ECG.

A second test involves administration of a compound as an infusion to pentobarbital anesthetized rats
25 in which the left ventricle is exposed to electrical square wave stimulation performed according to a preset protocol described in detail further below. This protocol includes the determination of thresholds for induction of extrasystoles and ventricular fibrillation. In addition,
30 effects on electrical refractoriness are assessed by a single extra beat technique. In addition effects on blood pressure, heart rate and the ECG are recorded. In this test, sodium channel blockers produce the ECG changes expected from the first test. In addition, sodium channel
35 blockers also raise the thresholds for induction of extrasystoles and ventricular fibrillation. Potassium

channel blockade is revealed by increasing refractoriness and widening of the Q-T intervals of the ECG.

A third test involves exposing isolated rat hearts to increasing concentrations of a compound. 5 Ventricular pressures, heart rate and ECG are recorded in the isolated heart in the presence of varying concentrations of the compound. This test provides evidence for direct toxic effects on the myocardium. Concentrations found to be effective in this test are 10 expected to be efficacious in the electrophysiological studies.

A fourth test is estimation of the antiarrhythmic activity of a compound against the arrhythmias induced by coronary artery occlusion in 15 anaesthetized rats. It is expected that a good antiarrhythmic compound will have antiarrhythmic activity at doses which have minimal effects on either the ECG, blood pressure or heart rate.

A compound is also tested directly for effects 20 on sodium and potassium currents in isolated rat myocytes. Isolated rat myocytes are obtained in the conventional manner used to obtain isolated myocytes from isolated hearts. They are used in conventional voltage clamp studies. In order to obtain adequate voltage clamps for 25 estimation of a compound's effects on sodium and potassium currents, the whole-cell patch clamp technique is used. In this technique, a microelectrode is attached to a cell in such a manner that the cell's internal contents are in free communication with the electrode's content. Using 30 the appropriate buffers and conventional voltage step protocols, both sodium and potassium currents can be identified. The effects of a compound are tested on these currents.

All of the foregoing tests are performed using 35 rat tissue. In order to ensure that a compound is not having effects which are only specific to rat tissue, further experiments are performed in dogs and primates.

In order to assess possible sodium channel and potassium channel blocking actions *in vivo* in dogs, a compound is tested for effects on the ECG, ventricular epicardial conduction velocity and responses to electrical stimulation. An anesthetized dog is subjected to an open chest procedure to expose the left ventricular epicardium. After the pericardium is removed from the heart a recording/stimulation electrode is sewn onto the epicardial surface of the left ventricle. Using this array, and suitable stimulation protocols, conduction velocity across the epicardium as well as responsiveness to electrical stimulation can be assessed. This information coupled with measurements of the ECG allows one to assess whether sodium and/or potassium channel blockade occurs. As in the first test in rats, a compound is given as a series of increasing bolus doses. At the same time possible toxic effects of a compound on the dog's cardiovascular system is assessed.

The effects of a compound on the ECG and responses to electrical stimulation are also assessed in intact halothane anesthetized baboons (*Papio anubis*). In this preparation, a blood pressure cannula and ECG electrodes are suitably placed in an anesthetized baboon. In addition, a stimulating electrode is placed into the right ventricle, together with a monophasic action potential electrode. As in the tests described above, ECG and electrical stimulation responses to a compound reveal the possible presence of sodium and/or potassium channel blockade. The monophasic action potential also reveals whether a compound widens the action potential, an action expected of a potassium channel blocker.

The present invention may be employed to treat the rhythm of a heart or prevent arrhythmias occurring in a heart that is susceptible to arrhythmia. Methods of administering effective amounts of antiarrhythmic agents are well known in the art and include the administration of an oral or parenteral dosage form. Such dosage forms

include, but are not limited to, parenteral solutions, tablets, capsules, sustained release implants, and transdermal delivery systems. Generally, oral or intravenous administration is preferred. The dosage amount and frequency will be selected to create an effective level of the agent without harmful effects. It will generally range from a dosage of from about 0.1 to about 100 mg/kg/day, and typically from about 0.1 to 10 mg/kg where administered orally or intravenously for antiarrhythmic effect. A 0.1% to 1% solution injected into a local site is typical for use as a local anaesthetic.

When the present invention is employed to induce local anaesthesia, the means of administration may be the same as described above in the case of treatment of arrhythmia, except that use of oral administration in the form of tablets or capsules will generally not be appropriate. Topical application of the local anaesthetic agent, for example in the form of an ointment or an aerosol spray, may be employed. Means of administering local anaesthetics are well known in the art.

Administration of this invention may be carried out in combination with the administration of other agents. For example, it may be desired to administer an opioid antagonist, such as naloxone, if a compound of formula I administered according to the present invention has not been selected from those enantiomers having reduced kappa opioid activity. The naloxone may antagonize opioid activity of the administered compound without adverse interference with the antiarrhythmic activity.

The present invention also includes a commercial kit containing a pharmaceutical composition which includes one or more compounds of formula I or, pharmaceutically acceptable salts thereof, in addition to any desired, pharmaceutically acceptable, carriers or diluents. The commercial kit also includes instructions for the use of

the pharmaceutical composition for the treatment of arrhythmia or for the inducement of local anaesthesia. Preferably the commercial package will contain one or more unit doses of the pharmaceutical composition. For
5 example, such a unit dose may be an amount sufficient for the preparation of an intravenous injection. It will be evident to those in the art that compounds which are light and/or air sensitive may require special packaging and/or formulation. For example, packaging may be used which is
10 opaque to light, and/or sealed from contact with ambient air, and/or formulated with suitable coatings or excipients.

The following examples are offered by way of illustration and not by way of limitation.

EXAMPLES

In the following examples, reference will be made to tests of a compound which is identified by name and the source of the compound tested, as follows:

U-50,488H:

trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]-3,4-dichlorophenylacetamide;

UpJohn Company.

In the following examples, antiarrhythmic efficacy was assessed against coronary occlusion induced arrhythmias in the rat using standard methods. ECG and electrophysiological stimulating tests in rats and primates involved standard ECG leads and ventricular stimulation (implanted electrodes in the rats and inserted right ventricular electrodes in primates). Intracellular recordings were made from the epicardium of rat hearts *in vivo*. Whole cell voltage clamp studies were performed in single, isolated, adult rat heart cells.

In the above array of tests, the compounds for use in this invention were typically found to be more than ten times as potent as some other Class I antiarrhythmics such as mexilitine (Igwezie, et al. (1992) *European Journal of Pharmacology*, 210:271-277), disopyramide, procainamide, and quinacainol (Penz, et al. (1992) *Journal of Pharmacological Methods*, 27:51-58). The compounds for use in this invention were equipotent with the extremely toxic tetrodotoxin (Abraham, et al. (1990) *J. Pharmacol. Exp. Ther.*, 251:1166-1173), which is a potent blocker of the cardiac sodium channel. However, the compounds for use in this invention did not exhibit the neuronal toxicity of tetrodotoxin or even that of lidocaine. No overt signs of anaesthesia resulted when the compounds were injected intravenously at antiarrhythmic doses.

Thus, the present invention may be used to treat arrhythmia, for example, by injection of the compound, without producing anaesthesia. However, the compounds may also be administered locally to induce a local anaesthetic effect.

It is estimated that the LD₅₀ of compound 10 in rats is 16 µmol/kg. For compound 1, the estimated LD₅₀ values in rats is 32 µmol/kg and in mice is 27 µmol/kg. Toxic symptoms of doses exceeding what is required for antiarrhythmic activity include fall in blood pressure or heart rate, and alteration of asystole/idioventricular rhythms. The bradycardic and hypotensive effects of the compounds suitable for this invention seem to be consistent with type I antiarrhythmic compounds but, the neuronal toxicity appears to be less than that of compounds such as lidocaine.

EXAMPLE 1

20

trans-N-methyl-2-(1-pyrrolidinyl)cyclohexanamine

This compound was prepared based on the method in U.S. Patent No. 4,579,863.

25

(i) Cyclohexene oxide (202 mL, 2 mol) was added dropwise to aqueous methyl amine (466 mL or 40% solution, 6 mol) over 70 minutes. After a further 90 minutes, the temperature of the reaction mixture was 48°C and was reduced to 30°C by cooling in a water bath. After a further 2 hours, the mixture had returned to room temperature. It was stirred overnight, and then refluxed for 3 h. The mixture was saturated with sodium hydroxide (cooled during addition), extracted several times with diethyl ether (total 500 mL), the diethyl ether layer dried over sodium sulphate overnight, and the diethyl ether removed on a rotary evaporator. The remaining

diethyl ether and cyclohexene oxide were removed by partial vacuum distillation. Distillation under full vacuum yielded a colorless fraction boiling at 95°C, (±)-trans-2-(methylamino)cyclohexanol: 217 g (84%).

- 5 (ii) A mixture of (±)-trans-2-(methylamino)cyclohexanol (200 g, 1.55 mol) and diethyl ether (400 mL) in a 3 L beaker was stirred and cooled in an ice bath as chlorosulfonic acid (103 mL, 1.55 mol) was
10 added dropwise. After approximately 25 mL had been added, it was necessary to stir the thick mixture with a spatula, and after a further 40 mL of acid had been added more diethyl ether (200 mL) was added. The whole addition took 105 minutes. The sticky mixture was stirred by hand and
15 left at room temperature for 2.5 hours. The mixture was filtered, and the solid washed with diethyl ether (300 mL). A solution of sodium hydroxide pellets (216 g) in water (1 L) was cooled in an ice bath, and then added slowly to the cooled solid. The mixture became less
20 viscous and the addition was complete within 20 minutes. The mixture was left to stand overnight, then poured into a 2 L flask and steam distilled, with water being added from a dropping funnel to maintain constant volume in the distillation pot. After the diethyl ether had distilled,
25 an organic product co-distilled with the water at a head temperature of 92-100°C (600 mL of a 2-phase colorless mixture was collected), to leave a small quantity of dark amber material on the surface of the water remaining in the distillation pot. The distillate was saturated with
30 sodium hydroxide and extracted with diethyl ether (8 x 100 mL), the diethyl ether layer dried over sodium sulphate and the diethyl ether removed on a rotary evaporator to leave crude product (133 g) which was distilled under reduced pressure to give 7-methyl-7-
35 azabicyclo[4.1.0]heptane (77.9 g, 43%).

(iii) A solution of ammonium chloride (1.6 g) in water (100 mL) was added to 7-methyl-7-azabicyclo[4.1.0]heptane (70 g, 0.59 mol) under nitrogen. Pyrrolidine (210 mL, 2.5 mol) was added and the mixture was stirred and refluxed under nitrogen for 20 hours. Sodium hydroxide was added to saturate the aqueous phase and the mixture was extracted with diethyl ether (7 x 100 mL). The combined organic extracts were washed with water (2 x 10 mL), and dried over sodium sulphate. The diethyl ether was removed on a rotary evaporator and excess pyrrolidine (60 mL) was distilled off under low vacuum. The product, (\pm)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine, was distilled under full vacuum (46-48°C). Yield 86 g (77%).

EXAMPLE 2

(\pm)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide monohydrochloride

This compound was prepared according to the procedure described by C.R. Clark et al. in *J. Med. Chem.* 31:831-836, 1988. A solution of 4-thianaphtheneacetyl chloride (prepared by refluxing 4-thianaphtheneacetic acid (1.94 g, 10 mmol) with excess thionyl chloride) in dichloromethane (10 mL) was added dropwise to a solution of (\pm)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine prepared in Example 1 (1.84 g, 10 mmol) in dichloromethane (10 mL) at 0°C. After stirring at room temperature for 10 minutes, diethyl ether was added until no further precipitate resulted. The crude product was collected by filtration, washed with diethyl ether and dried in vacuo. It was recrystallised from methanol/diethyl ether, to give the title compound, 3.3 g (85%). Proton and carbon-13 NMR data in accord.

Elemental analysis: Calcd. for $C_{21}H_{29}N_2OClS$: C 64.18, H 7.44%, N 7.13%; Found C 63.34, H 7.17, N 7.10.

5

EXAMPLE 3

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] (3,4-dichlorophenoxy)acetamide monohydrochloride

10 The title compound was prepared according to the method described in Example 2, using 3,4-dichlorophenoxyacetic acid (2.34 g, 11 mmol) and (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine prepared in Example 1 (2 g, 11 mmol). The crude product,
15 which started to precipitate before the addition of the diethyl ether, was recrystallised from hot dichloromethane. Yield 2.9 g (62%).

Elemental analysis: Calcd. for $C_{19}H_{27}N_2O_2Cl_3$. CH_2Cl_2 : C 47.41, H 5.77, N 5.53 Cl 34.98%; Found C 47.86,
20 H 5.79, N 5.53, Cl 35.96%.

EXAMPLE 4

25

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-3-acetamide monohydrochloride

The title compound was prepared according to the
30 method described in Example 2, using 3-thianaphtheneacetic acid (4.24 g, 22 mmol) and (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine prepared in Example 1 (3.85 g, 21 mmol). The crude product, which started to precipitate before the addition of the diethyl ether, was
35 recrystallised from hot methanol/diethyl ether to yield a white solid, 6.54 g (79%).

Elemental analysis: Calcd. for $C_{21}H_{29}N_2OClS$: C 64.18, H 7.44, N 7.13%; Found C 63.16, H 7.40, N 6.99%.

5

EXAMPLE 5

Resolution of (\pm)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine

10 The resolution is based on a modification of the method reported by C.R. Clark et al. in *J. Med. Chem.* 31:831-836, 1988. The (-)-enantiomer (R,R) of trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine prepared in Example 1 was separated by repeated fractional
15 crystallization of the 2,3-di-p-toluoyl-D-tartaric acid salts. A solution of (\pm)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine (16 g, 88 mmol) in boiling methanol (400 mL) was treated with a solution of 2,3-di-p-toluoyl-D-tartaric acid (35.6 g, 88 mmol) in boiling
20 methanol (400 mL). The solution was cooled to 30°C and the resulting solid was collected by filtration. The above process was repeated for two further portions of racemic diamine (16 g each), and the three batches of solid (total 101 g, $[\alpha]^{20}_D(CH_2Cl_2)$ -68° for free diamine)
25 which were isolated were combined and washed with boiling methanol (1 L). The washed solid (78 g, $[\alpha]^{20}_D(CH_2Cl_2)$ -82° for free diamine) was divided into batches (approx. 5 g each), and each batch was recrystallised from boiling methanol (600 mL). The first crops (total 33.9 g, $[\alpha]^{20}_D(CH_2Cl_2)$ -93.2° for free diamine) were filtered off after
30 allowing the methanol solutions to stand at room temperature for 3-4 hours. The second crops (total 9.25 g, $[\alpha]^{20}_D(CH_2Cl_2)$ -92.6° for free diamine) were isolated from the filtrate after standing in the fridge overnight.
35 The two crops were combined and converted to the free diamine by partitioning between dichloromethane and aqueous (20%) potassium hydroxide solution. The

dichloromethane layer was dried and evaporated, and the residue was partitioned between diethyl ether and aqueous (10%) potassium hydroxide solution. The diethyl ether layer was dried and evaporated to give the product, (1R, 2R)-(-)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine, as a pale yellow oil 14.3 g (60%); $[\alpha]^{20}_D$ (CH₂Cl₂) -92.9°.

EXAMPLE 6

10

(1R,2R)-(+)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide monohydrochloride

15

The title compound was prepared according to the method described in Example 2, using 4-thianaphtheneacetic acid (2.28 g, 12 mmol) and (1R,2R)-(-)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]amine isolated using the resolution method described in Example 5 (2.21 g, 12 mmol). The title compound (the (+)-enantiomer of the racemate prepared in Example 2) was recrystallised from hot methanol. Yield 3.26 g, (69%). $[\alpha]^{20}_D$ (CH₂Cl₂) +29.3°. Proton and carbon-13 NMR data in accord.

Elemental analysis: Calcd. for C₂₁H₂₉N₂OCls: C 64.18, H 7.44%, N 7.13%; Found C 61.49, H 7.45, N 6.72%.

25

EXAMPLE 7

30

[(±)-(1α,2β,4β,5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]amine

The starting material, [(±)-(1α,2β,4β,5β)]-[4,5-dimethoxy-2-(1-pyrrolidinyl)]cyclohexanol, was prepared according to the procedure described in U.S. Patent No. 4,855,316. This was converted to the title compound as follows:

35

A solution of [(±)-(1 α ,2 β ,4 β ,5 β)]-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexanol (7 g, 30.5 mmol) and triethylamine (4.8 mL, 35 mmol) in dichloromethane (70 mL) under nitrogen was cooled in an ice-bath as
5 methanesulfonyl chloride (2.55 mL, 33 mmol) in dichloromethane (35 mL) was added dropwise. After 1 hour at 0°C, GC showed the reaction to be complete. The mixture was diluted with dichloromethane (70 mL) and washed with water (2 x 50 mL). The aqueous phase was
10 washed with dichloromethane (20 mL) and the organic fractions were combined and dried over sodium sulphate. The solvent was removed in vacuo to leave a yellow oil. An aqueous solution of methylamine (50 mL of a 40% solution, 0.57 mol) was added to the mesylated product,
15 and the mixture refluxed for a total of 1.5 hours. The reaction mixture was partitioned between 10% sodium hydroxide solution (150 mL) and dichloromethane (150 mL). The aqueous layer was washed with dichloromethane (150 mL) and the combined organic layers were dried over sodium
20 sulphate. The solvent was removed to leave the product as a yellow oil. Yield 7 g (95%). Proton and carbon-13 NMR data in accord.

25

EXAMPLE 8

[(±)-(1 α ,2 β ,4 β ,5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide
monohydrochloride

30

The title compound was prepared according to the method described in Example 2, using 4-thianaphtheneacetic acid (0.78 g, 4 mmol) and [(±)-(1 α ,2 β ,4 β ,5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]amine prepared
35 in Example 7 (1 g, 4 mmol). The product was recrystallised from hot methanol/diethyl ether.

Elemental analysis: Calcd. for $C_{23}H_{33}N_2O_3ClS$: C 60.98, H 7.43, N 6.18; Found C 60.79, H 7.30, N 6.13%.

5

EXAMPLE 9

$[(\pm)-(1\alpha, 2\beta, 4\beta, 5\beta)]$ -N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide
monohydrochloride

10

The title compound was prepared according to the method described in Example 2, using 3,4-dichlorophenoxyacetic acid (0.91 g, 4 mmol) and $(\pm)-(1\alpha, 2\beta, 4\beta, 5\beta)]$ -N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]amine prepared in Example 7 (1 g, 4 mmol). The product was washed with diethyl ether. Yield 1.2 g (60%).

15

Elemental analysis: Calcd. for $C_{21}H_{31}N_2O_4Cl_3$: C 52.35, H 6.48, N 5.81%, Found C 52.15, H 6.52, N 5.75%.

20

EXAMPLE 10

(\pm) -trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl]amine

25

(i) Hexamethyleneimine (17.3 mL, 0.15 mol), cyclohexene oxide (15.5 mL, 0.15 mol) and water (5 mL) were refluxed under nitrogen overnight. GC analysis showed the reaction to be complete. The cooled mixture was partitioned between saturated sodium hydroxide solution (50 mL) and diethyl ether (75 mL). The aqueous layer was washed with diethyl ether (30 mL), and the combined diethyl ether layers were dried over sodium sulphate. The diethyl ether was removed from the amber solution on a rotary evaporator. The crude product, (\pm) -

30

35

trans-[2-(1-hexahydroazepinyl)]cyclohexanol (28 g), was purified by full vacuum distillation (62-65°C) 23 g (76%).

(ii) A solution of (\pm)-*trans*-[2-(1-hexahydroazepinyl)]cyclohexanol prepared above (17 g, 86 mmol) and triethylamine (13.5 mL, 97 mmol) in dichloromethane (135 mL) under nitrogen was cooled in an ice-bath as methanesulfonyl chloride (7.25 mL, 93 mmol) in dichloromethane (70 mL) was added dropwise. After 50 minutes at 0°C, GC showed the reaction to be complete. The mixture was diluted with dichloromethane (135 mL) and washed with water (2 x 135 mL). The aqueous phase was washed with dichloromethane (70 mL) and the organic fractions were combined and dried over sodium sulphate. The solvent was removed in vacuo to leave a yellow oil (19.7 g). An aqueous solution of methylamine (140 mL of a 40% solution, 1.6 mol) was added to the mesylated product, and the mixture refluxed for a total of 5.5 h. The reaction mixture was partitioned between 10% sodium hydroxide solution (335 mL) and dichloromethane (335 mL). The aqueous layer was washed with dichloromethane (200 mL) and the combined organic layers were dried over sodium sulphate. The solvent was removed to leave the crude product as a yellow oil. This was distilled under full vacuum (bp 103-105°C) to give the title diamine as a colorless liquid. Yield 15.5 g (86%).

EXAMPLE 11

(\pm)-*trans*-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide monohydrochloride

The title compound was prepared according to the method described in Example 2, using 3,4-dichlorophenoxyacetic acid (2.15 g, 10 mmol) and (\pm)-

trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl]amine prepared in Example 10 (2 g, 10 mmol). The crude product, which precipitated without the addition of any diethyl ether, was recrystallised from hot methanol (150 mL).

5 Yield 3.58 g (80%).

Elemental analysis: Calcd. for $C_{21}H_{31}N_2O_2Cl_3$: C 56.07, H 6.95, N 6.23 Cl 23.64%; Found C 55.29, H 7.20, N 6.02, Cl 23.84%.

10

EXAMPLE 12

(±)-*trans*-N-methyl-N-[2-(morpholinyl)cyclohexyl]amine

15 (i) The intermediate aminoalcohol, (±)-*trans*-[2-(morpholinyl)]cyclohexanol, was prepared by the method detailed in Example 10(i), refluxing morpholine (5 g, 57 mmol), cyclohexene oxide (5.8 mL, 57 mmol) and water (3 mL) for 1.5 h. The crude product (9.8 g) was purified by
20 full vacuum distillation (bp 75-80°C) 8.7 g (83%).

(ii) The aminoalcohol (4.6 g, 25 mmol) was converted to the title compound using the method detailed in Example 10(ii), using triethylamine (3.9 mL, 28 mmol)
25 and methanesulfonyl chloride (2.1 mL, 27 mmol), followed by treatment (1 hour reflux) with aqueous methylamine (50 mL of a 40% solution, 0.46 mol). The crude title product, which was not distilled, was obtained as a pale yellow oil which crystallized overnight. Yield 4.73 g (96%).

30

EXAMPLE 13

35

Antiarrhythmic efficacy was assessed by investigating the activity of the compounds on the

incidence of cardiac arrhythmias in pentobarbital anaesthetized rats subject to coronary artery occlusion. Rats weighing 150-200 gm were subjected to preparative surgery and assigned to groups in a random block design.

5 In each case, the trachea was cannulated and the animals were artificially ventilated. The left carotid artery was cannulated for measurement of mean arterial blood pressure and withdrawal of blood samples. The right jugular vein was also cannulated for injection of drugs. The thoracic

10 cavity was opened and a polyethylene occluder loosely placed around the left anterior descending coronary artery. ECG was recorded by insertion of electrodes placed along the anatomical axis of the heart determined by palpation. In random and double-blind manner, rats

15 were given an initial injection of saline or, 8 μ mol/kg naloxone hydrochloride (Dupont Pharmaceutical Co.). A second injection of saline vehicle, or the compound to be tested at various doses, was given 5 minutes later. All drugs were dissolved in 0.9% NaCl solution. Thereafter,

20 the occluder was pulled so as to produce coronary artery occlusion. ECG, arrhythmias, blood pressure, heart rate on mortality were monitored for 30 minutes after occlusion. Arrhythmias were recorded as ventricular tachycardia (VT) and ventricular fibrillation (VF) and

25 according to Curtis, M.J. and Walker, M.J.A. (1988) *Cardiovasc. Res.*, 22:656. Rats were excluded from the study if they did not exhibit pre-occlusion serum potassium concentrations within the range of 2.9 - 3.9 mM; occlusion associated with increases in R-wave height and

30 "S-T" segment elevation; and, an occluded zone (measured after death by cardiogreen dye perfusion) in the range of 25-50% of total left-ventricular weight.

Table 1 describes the result of tests of the compounds described therein as ED₅₀ values which are the

35 doses required to produce 50% reductions in the arrhythmic activity referred to therein. Initial results with compound 7 showed that at 2 and 8 μ mol/kg, the incidents

of fatal arrhythmias (VF) was reduced to 25% and 0%, respectively, from a control value of 88%. Naloxone alone slightly decreased the incidence of VF arrhythmias. Pretreatment with naloxone did not antagonize the activity of the compounds shown in Table 1. The doses of U-50,488H and compound 7 required for significant antiarrhythmic effects exceeded the doses required for kappa agonism for those compounds. These results demonstrate that the compounds useful within this invention are significantly more potent antiarrhythmic agents than U-50,488H, especially in respect of fatal arrhythmias (VF). Therefore, the former compounds have a much more useful therapeutic ratio (which compares efficacy to toxicity) than U-50,488H.

TABLE 1

	<u>Activity</u>	<u>Compound 10</u>	<u>Compound 7</u>	<u>U-50,488H</u>
20	VT	6	8	16
	VF	1	<1	6

EXAMPLE 14

25 The procedures described in Example 13 were carried out and measurements taken before coronary artery occlusion. Table 2 describes blood pressure (BP), heart rate (HR), and EKG (PR; QRS; and, QT intervals) effects of compound 7 at 2 and 8 $\mu\text{mol/kg}$ in the presence and absence of naloxone (8 $\mu\text{mol/kg}$). Values are the mean \pm sem in pentobarbital anaesthetized rats (n=8), and * indicates $P < 0.05$ of differences from control. Measurements were made 10 minutes after the end of drug administration.

35 Compound 7 prolonged P-R interval and QRS width as well as depressing blood pressure and heart rate in a manner generally unaffected by naloxone. This is evidence

that the compound causes sodium channel blockade in a manner independent of its kappa opioid agonist activity.

5

TABLE 2

<u>Treatment</u>						
Initial:	saline	naloxone	saline	saline	naloxone	naloxone
Final:	saline	saline	Cmpd. 7	Cmpd. 7	Cmpd. 7	Cmpd. 7
Dose of Cmpd. 7	-	-	2.0	8.0	2.0	8.0
BP	109±5	103±5	88±4*	88±5*	75±5*	80±3*
HR	377±12	379±6	356±16*	331±11*	298±9*	353±12
PR	53±1	57±1*	53±2	62±2*	57±2	58±1*
QRS	28±1	29±1	31±1*	32±1*	33±1*	32±1*
QT	38±1	41±1	37±1	42±0.5*	41±1	41±0.5*

10

EXAMPLE 15

The procedures of Example 14 were carried out in respect of various doses of compound 10, compound 7, and U-50,488H. Table 3 describes the results of the tests as

15 ED₂₀ (μmol/kg) which are the doses required to produce a 20% change in the indice measured. The increases in P-R interval and QRS interval indicate cardiac sodium channel blockade while the increase in Q-T interval indicates ancillary cardiac potassium channel blockade which is the

20 property of a type Ia antiarrhythmic. The electrophysiological effects of compound 10 and compound 7 occur

very rapidly, generally achieving a maximum effect in one circulation time.

Similar results have been obtained in primates, wherein compound 1 appears to be equipotent to compound 10 and compound 7.

TABLE 3

		Cmpd. 10	Cmpd. 7	U-50,488H
EKG P-R	interval increase	10	16	16
EKG QRS	interval increase	>32	>32	32
EKG Q-T	interval increase	32	>32	32

10

EXAMPLE 16

Rats were prepared according to the preceding procedures, but without the coronary artery occlusion. Two Teflon coated silver wires stimulating electrodes were inserted through the chest wall and implanted in the left ventricle. Square wave stimulation was used to determine threshold current for capture, ventricular fibrillation threshold current, and effective refractory period (Howard, P.G. and Walker, M.J.A. (1990) *Proc. West. Pharmacol. Soc.*, 33:123-127). Intracellular action potential duration and rise rate was determined according to the methods described in Abraham, et al., (1990) *J. Pharmacol. Exp. Ther.*, 251:1166-1173. Table 4 contains ED₂₀ values for these indices of cardiac sodium channel blockage in respect of the named compounds. The increases in refractoriness and action potential duration indicate ancillary blockade of potassium channels. The effect on action potential rise rate by compound 1 is indicative of cardiac sodium channel blockade.

TABLE 4

		Threshold current increase	Fibrillation current increase	Refractoriness increase
5	Compound			
	1	0.3	0.3	0.3
	2	0.2	0.2	0.4
10	3	0.2	0.2	0.4
	4	0.4	0.7	1
	5	1	0.3	0.6
	7	8	8	10
	8	2	1	2
15	9	0.3	0.3	0.3
	10	1	1	10
	U-50, 488H	16	16	10

20

EXAMPLE 17

In order to directly measure cardiac sodium channel blockade, standard whole cell voltage clamp tests were carried out using single isolated adult rat cardiac cells. The tests were carried out in the presence of naloxone which blocked kappa opioid activity of the compounds being tested. Results in the form of EC₂₀ values, which are micromolar concentrations required to produce a 20% change in activity, are presented in Table 5. These results show that compound 1 is a potent sodium channel blocker like compound 7. Similar whole cell voltage clamp analysis with compound 1 demonstrated an effect on potassium current indicative of blockade at an EC₂₀ of 100 μ molar.

TABLE 5

	<u>Compound</u>	<u>Sodium current reduction</u>
5	1	4
	2	2
	6	13
	7	10
	8	4
10	9	4


EXAMPLE 18

15 The guinea pig intradermal wheal assay for local
anaesthesia was carried out. The test consisted of
injecting intradermally a small volume of approximately
0.1 ml of the test compound at various concentrations in a
saline vehicle into the back of guinea pig, after which
20 local anaesthesia was determined by the absence of a
flinch upon pin prick near the site of injection. In this
test, ED₂₀ values for compound 1; compound 7; and
U-50,488H were 0.3, 0.5, and >0.5 percent (g/100 ml),
respectively.


25 From the foregoing, it will be evident that,
although specific embodiments of the invention have been
described herein for purposes of illustration, various
modifications may be made without deviating from the
30 spirit and scope of the invention.

Claims


1. An enantiomer or geometric isomer of a compound of formula I, or a pharmaceutically acceptable salt thereof, for use within a method of blocking sodium channels in cardiac or neuronal tissue, said compound of the formula:



wherein n is either 0 or 1; R₁, R₂, R₃, R₄ are hydrogen, hydroxy, alkoxy of from one to four carbon atoms, or points of attachment of a spiro- or fused five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; R₅ and R₆ are either alkyl of from one to five carbon atoms or, when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl, piperidinyl, or hexahydroazepinyl ring; and Q is selected from the group of substituents comprising: 3,4,5-trimethylphenoxy;




where R₇ is hydrogen, fluorine, chlorine, alkyl of from one to six carbon atoms, or aryl; Z is -CH₂-, -O-, -S-, or N-R₈ where R₈ is hydrogen, alkanoyl of from one to six carbon atoms, or alkyl of from one to six carbon atoms;




(III)

where R_9 and R_{10} are independently hydrogen, fluorine, bromine, alkyl of from one to six carbon atoms, or alkoxy of from one to four carbon atoms;



(IV)

where R_9 and R_{10} are defined as above; and



(V)

where R_{11} and R_{12} are independently hydrogen, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl.

2. A compound according to claim 1 wherein the stereoisomeric arrangement of the amine and amide bearing carbon atoms of the cyclohexyl ring of the compound formula I are as depicted in claim 1.

3. A compound according to claim 1 wherein $n=1$; R_5 and R_6 taken together with the nitrogen atom to which they are attached form a pyrrolidinyl ring; R_3 and R_4 are hydrogen; R_1 and R_2 are selected from the group comprising hydrogen and an

oxaspiran ring; and Q is selected from the group comprising substituents II, III, and IV.

4. A compound according to claim 3 wherein Q is substituent II.

5. A compound according to claim 2 wherein $n=1$; R_5 and R_6 taken together with the nitrogen atom to which they are attached form a pyrrolidinyl ring; R_3 and R_4 are hydrogen; R_1 and R_2 are selected from the group comprising hydrogen and points of attachment of an oxaspiran ring; and Q is selected from the group comprising substituents II, III and IV.

6. A compound according to claim 5 wherein Q is substituent II.

7. A compound according to claim 6 wherein the method is for arrhythmia by blocking sodium channels in cardiac tissue.

8. A compound according to claim 1 wherein the compound is selected from the group consisting of:

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] benzo[b]thiophene-4-acetamide;

(1R,2R) - (+) -N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] benzo[b]thiophene-4-acetamide;

[(±)-(1 α ,2 β ,4 β ,5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;

[(±)-(1 α ,2 β ,4 β ,5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy) acetamide;

(±)-trans-N-methyl-N-[2-(1-hexahydroazepinyl) cyclohexyl](3,4-dichlorophenoxy) acetamide;

(±)-trans-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;

[5R(5 α ,7 α ,8 β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] (3,4-dichlorophenoxy) acetamide;

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] benzo[b]thiophene-3-acetamide;

[5S(5α, 7α, 8β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;

(1S, 2S)-2-(benzo[b]thiophen-4-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1R, 2R)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1S, 2S)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1R, 2R)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1S, 2S)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1R, 2R)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1S, 2S)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

[1S(1α, 2β, 4β)]-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide;

[1R(1α, 2β, 4β)]-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide;

(1R, 2R)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide; and

(1S, 2S)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide.

9. A compound according to claim 1 wherein the compound is selected from the group consisting of

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] benzo[b]thiophene-4-acetamide;

(1R, 2R) - (+) -N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl] benzo[b]thiophene-4-acetamide;

[(±)-(1α, 2β, 4β, 5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;

[(±)-(1α, 2β, 4β, 5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl] (3,4-dichlorophenoxy) acetamide; and

(±)-trans-N-methyl-N-[2-(1-hexahydroazepinyl) cyclohexyl] (3,4-dichlorophenoxy) acetamide.

10. A compound according to claim 1 wherein the compound is selected from the group consisting of:

[5S(5 α , 7 α , 8 β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-benzo[b]furan-4-acetamide;
(1R,2R)-2-(benzo[b]thiophen-4-yl)-N-methyl-N-[2-(1-pyrrolidinyl) cyclohexyl] acetamide;
(1R,2R)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl) cyclohexyl] acetamide;
(1R,2R)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl) cyclohexyl] acetamide;
(1R,2R)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl) cyclohexyl] acetamide;
[1R(5 α , 2 β , 4 β)]-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl) cyclohexyl] benzo[b]furan-4-acetamide; and
(1R,2R)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino) cyclohexyl] carboxamide.

11. A compound according to claim 1 wherein the compound is:

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl) cyclohexyl] benzo[b]thiophene-4-acetamide.

12. A compound according to claim 1 wherein the compound of formula I is:

(1R,2R)-(+)-N-methyl-N-[2-(1-pyrrolidinyl) cyclohexyl] benzo[b]thiophene-4-acetamide.

13. A compound according to claim 1 wherein the compound is:

[(±)-(1 α , 2 β , 4 β , 5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl) cyclohexyl] benzo[b]thiophene-4-acetamide.


14. A compound according to claim 1 wherein the compound is:

$[(\pm)-(1\alpha, 2\beta, 4\beta, 5\beta)]$ -N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy) acetamide.

15. A compound according to claim 1 wherein the compound is:


(\pm) -trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide.

16. Use of an enantiomer or geometric isomer of a compound of formula I, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use in blocking sodium channels in cardiac or neuronal tissue, said compound of the formula:




(I)

wherein n is either 0 or 1; R₁, R₂, R₃, R₄ are hydrogen, hydroxy, alkoxy of from one to four carbon atoms, or points of attachment of a spiro- or fused five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; R₅ and R₆ are either alkyl of from one to five carbon atoms or, when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl, piperidinyl, or hexahydroazepinyl ring; and Q is selected from the group of substituents comprising: 3,4,5-trimethylphenoxy;




(II)

where R_7 is hydrogen, fluorine, chlorine, alkyl of from one to six carbon atoms, or aryl; Z is $-CH_2-$, $-O-$, $-S-$, or $N-R_8$ where R_8 is hydrogen, alkanoyl of from one to six carbon atoms, or alkyl of from one to six carbon atoms;




(III)

where R_9 and R_{10} are independently hydrogen, fluorine, bromine, alkyl of from one to six carbon atoms, or alkoxy of from one to four carbon atoms;



(IV)

where R_9 and R_{10} are defined as above; and



(V)

where R_{11} and R_{12} are independently hydrogen, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl.

17. Use of a compound for the manufacture of a medicament for use in blocking sodium channels in cardiac or neuronal tissue, said compound selected from the group consisting of:

(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
(1R,2R)-(+)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
[(±)-(1α,2β,4β,5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
[(±)-(1α,2β,4β,5β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
(±)-trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
(±)-trans-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;
[5R(5α,7α,8β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;
(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
(±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-3-acetamide;
[5S(5α,7α,8β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzo[b]furan-4-acetamide;
(1S,2S)-2-(benzo[b]thiophen-4-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
(1R,2R)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
(1S,2S)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
(1R,2R)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
(1S,2S)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
(1R,2R)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
(1S,2S)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

[1S(1 α , 2 β , 4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide;
 [1R(1 α , 2 β , 4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide;
 (1R, 2R)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide; and
 (1S, 2S)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide.

18. Use of a compound for the manufacture of a medicament for use in blocking sodium channels in cardiac or neuronal tissue, said compound selected from the group consisting of:

(\pm)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
 (1R, 2R)-(+)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
 [(\pm)-(1 α , 2 β , 4 β , 5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]thiophene-4-acetamide;
 [(\pm)-(1 α , 2 β , 4 β , 5 β)]-N-methyl-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide;
 and
 (\pm)-trans-N-methyl-N-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetamide.

19. Use of a compound for the manufacture of a medicament for use in blocking sodium channels in cardiac or neuronal tissue, said compound selected from the group consisting of:

[5S(5 α , 7 α , 8 β)]-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-benzo[b]furan-4-acetamide;
 (1R, 2R)-2-(benzo[b]thiophen-4-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
 (1R, 2R)-2-(indol-3-yl)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
 (1R, 2R)-2-(2,3-dichlorophenoxy)-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

(1R,2R)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;

[1R(5 α ,2 β ,4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]benzo[b]furan-4-acetamide; and

(1R,2R)-inden-2-yl-N-methyl-N-[2-(1,1-dimethylamino)cyclohexyl]carboxamide.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 93/00124

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 C07D295/12; C07D333/60; C07D333/54; A61K31/16 A61K31/33; A61K31/34; A61K31/38		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07D ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	EP,A,0 146 297 (WARNER-LAMBERT CO. ; USA (US)) 26 June 1985 cited in the application see page 7, line 22 - page 8, line 29 see examples ---	1-10, 16-19
X	EP,A,0 147 085 (WARNER-LAMBERT CO.; USA (US)) 3 July 1985 cited in the application see page 8, line 18 - page 9, line 28 see page 15 see examples --- <div style="text-align: right;">-/--</div>	1-12, 16-19
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but, later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search <div style="text-align: center;">26 JULY 1993</div>	Date of Mailing of this International Search Report <div style="text-align: center;">30. 07. 93</div>	
International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center;">PAUWELS G.R.A.</div>	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	US,A,4 663 343 (WARNER-LAMBERT CO.; USA (US)) 5 May 1987 cited in the application see column 4, line 14 - line 50 see claims; examples ---	1-5,8, 16,17
X	EP,A,0 207 773 (WARNER-LAMBERT CO.; USA (US)) 7 January 1987 cited in the application see page 10, line 17 - page 15, line 20 see examples ---	1-8,10, 16,17,19
X	US,A,4 855 316 (WARNER-LAMBERT CO.; USA (US)) 8 August 1989 cited in the application see column 4, line 41 - column 6, line 45 see examples ---	1-8,9, 13,16-18
X	EP,A,0 372 466 (WARNER-LAMBERT CO.; USA (US)) 13 June 1990 see page 5, line 36 - line 41 see examples ---	1-8,10, 16,17,19
A	CHEMICAL ABSTRACTS, vol. 117 Columbus, Ohio, US; abstract no. 48285, MA S C ET AL. 'Synthesis and analgesic activity of analogs of U-50488, an opiate kappa-agonist' see abstract & YAOXUE XUEBAO (YHHPAL,05134870); 91; VOL.26 (12); PP.902-5 ---	1
X	J. MED. CHEM. ; 88; VOL.31 (4); PP.831-6 'Highly selective .kappa. opioid analgesics. Synthesis and structure-activity relationships of novel N-[(2-aminocyclohexyl)aryl] acetamide and N-[(2-aminocyclohexyl)aryloxy]acetamide derivatives' cited in the application see page 832; table I ---	1-13, 16-19
	---	-/--

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	J. MED. CHEM. ; 89; VOL.32 (7); PP.1620-6 'Highly selective .kappa.-opioid analgesics. 2. Synthesis and structure activity relationships of novel N-(2-aminocyclohexyl)arylac etamide derivatives' * page 1621, Scheme I, compounds 5-12 *	1-8, 16, 17
X	J. MED. CHEM. ; 90; VOL.33 (1); PP.286-91 'Highly selective .kappa.-opioid analgesics. 3. Synthesis and structure-activity relationships of novel N-[2-(1-pyrrolidinyl)-4- or -5-substituted cyclohexyl]arylacetamide derivatives' cited in the application see page 288; table II see page 289; table III	1-8, 10, 16, 17, 19
X	EP,A,0 380 063 (WARNER-LAMBERT COMP.) 1 August 1990 see page 2, line 35 - page 4, line 21 see page 7, line 1	16
X	WO,A,8 607 257 (THE UPJOHN COMPANY) 18 December 1986 cited in the application see page 4, line 10 - page 5, line 27	16

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

CA 9300124
SA 72870

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 26/07/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0146297	26-06-85	US-A- 4598087	01-07-86
		AU-B- 565237	10-09-87
		AU-A- 3630684	13-06-85
		CA-A- 1235122	12-04-88
		DE-A- 3469168	10-03-88
		JP-A- 60166647	29-08-85
		OA-A- 7884	20-11-86
		US-A- 4677122	30-06-87
EP-A-0147085	03-07-85	US-A- 4656182	07-04-87
		AU-B- 565236	10-09-87
		AU-A- 3608084	13-06-85
		CA-A- 1234808	05-04-88
		JP-A- 60166662	29-08-85
		OA-A- 7885	20-11-86
		US-A- 4738981	19-04-88
US-A-4663343	05-05-87	None	
EP-A-0207773	07-01-87	US-A- 4737493	12-04-88
		AU-B- 578837	03-11-88
		AU-A- 5919486	08-01-87
		CA-A- 1264321	09-01-90
		JP-A- 62059271	14-03-87
US-A-4855316	08-08-89	None	
EP-A-0372466	13-06-90	AU-B- 626949	13-08-92
		AU-A- 4586189	14-06-90
		CA-A- 2004575	06-06-90
		JP-A- 2200659	08-08-90
		US-A- 5051428	24-09-91
EP-A-0380063	01-08-90	US-A- 4906655	06-03-90
		AU-B- 630411	29-10-92
		AU-A- 4869990	02-08-90
		CA-A- 2008391	24-07-90
		JP-A- 2233669	17-09-90
		US-A- 5019588	28-05-91

EPO FORM P067

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

CA 9300124
SA 72870

26/07/93
2

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-8607257	18-12-86	EP-A- 0263208	13-04-88
		EP-A- 0224566	10-06-87
		JP-T- 63500796	24-03-88
